

P1.1

P1.1

$$\int_{\frac{1}{2}}^2 (e^x - 1) dx \quad \left| \begin{array}{l} e^x = y \\ e^x dx = dy \\ dx = \frac{dy}{y} \\ y = 2 \end{array} \right. \quad = \int_{\frac{1}{2}}^2 \frac{\sqrt{y-1}}{2} dy = \left| \begin{array}{l} t = \sqrt{y-1} \\ dt = \frac{dy}{2\sqrt{y-1}} \end{array} \right. =$$

$$e^0 = 1$$

$$e^{2^2} = 2$$

$$dt dt = dy$$

$$\text{LB: } t=0$$

$$\text{UB: } t=\sqrt{2}$$

$$t^2 = y-1$$

$$t^2 = y^2$$

$$t = y^2$$

$$t^2 = y-1$$

$$= 2 \int_0^1 \frac{t^2}{t^2 + 1} dt = 2 \int_0^1 \left(1 - \frac{1}{1+t^2} \right) dt$$

$$= 2t \Big|_0^1 - 2 \arctan t \Big|_0^1 = 2(1-0) - 2 \left(\frac{\pi}{4} - 0 \right) = \underline{\underline{2 - \frac{\pi}{2}}}$$

P1.2

$$\int \arccos x = \left| \begin{array}{l} u = \arccos x \quad u' = \frac{-1}{\sqrt{1-x^2}} \\ v' = 1 \quad v = x \end{array} \right| = \int x \arccos x \Big|_0^1$$

$$+ \int_0^1 \frac{x}{\sqrt{1-x^2}} dx = -0 \cdot \frac{\pi}{2} - 1(0) - \sqrt{1-x^2} \Big|_0^1 = \underline{\underline{1}}$$

$$\cancel{2(\sqrt{1-x^2})}$$

P1.3

$$\int x^{2k-1} e^{-\frac{x^2}{2}} \quad (k \in \mathbb{N} \quad (\text{nat. num.}))$$

P1.3

$$\int_0^\infty x^{2k-1} e^{-\frac{x^2}{2}} dx = \begin{cases} \frac{x^2}{2} = t \\ xdx = dt \end{cases}$$

(P1-12)

$$= \int_0^\infty (2t)^{k-1} e^{-t} dt$$

$$= \int_0^\infty x^{2k-2} e^{-\frac{x^2}{2}} \underbrace{x dx}_{(x^2)(k-1)} \underbrace{dt}_{e^{-t}}$$

$$\begin{aligned} x^2 &= 2t \\ x &= \sqrt{2t} \end{aligned}$$

$$= \int_0^\infty (2t)^{k-1} e^{-t} dt$$

$$= 2^{k-1} \int_0^\infty t^{k-1} e^{-t} dt$$

$\equiv F(k)$

$F(z) + z \in \mathbb{C} \setminus \{\text{non-positive integers}\}$

$F(n) = (n-1)! \text{ for } n \in \mathbb{N}$

$$= 2^{k-1} (k-1)!$$

$$\begin{aligned} \int_0^\infty t^{k-1} e^{-t} dt &\stackrel{?}{=} \left[-t^{k-1} e^{-t} \right]_0^\infty + (k-1) \int_0^\infty t^{k-2} e^{-t} dt \\ &\stackrel{?}{=} (k-1) \left[-t^{k-2} e^{-t} \right]_0^\infty + (k-1)(k-2) \int_0^\infty t^{k-3} e^{-t} dt \end{aligned}$$

$$\begin{aligned} &= (k-1)(k-2) \cdots 2 \cdot 1 \int_0^\infty t^0 e^{-t} dt = (k-1)! \int_0^\infty e^{-t} dt \\ &= (k-1)! \left[-e^{-t} \right]_0^\infty = (k-1)! \end{aligned}$$

Algebra

Applied Mathematics

Calculus and Analysis

Discrete Mathematics

Foundations of Mathematics

Geometry

History and Terminology

Number Theory

Probability and Statistics

Recreational Mathematics

Topology

Alphabetical Index

Interactive Entries

Random Entry

New in MathWorld

MathWorld Classroom

About MathWorld

Contribute to MathWorld

Send a Message to the Team

MathWorld Book

Wolfram Web Resources

13,612 entries

Last updated: Mon Feb 27 2017


Created, developed, and
nurtured by Eric Weisstein
at Wolfram Research

Calculus and Analysis > Special Functions > Gamma Functions >
 Calculus and Analysis > Special Functions > Named Integrals >
 Calculus and Analysis > Special Functions > Product Functions >
 More...

Gamma Function

DOWNLOAD
Wolfram Notebook

VIEW | DOWNLOAD | THE MATHEMATICA NOTEBOOK



The (complete) gamma function $\Gamma(z)$ is defined to be an extension of the factorial to complex and real number arguments. It is related to the factorial by

$$\Gamma(n) = (n-1)!.$$

gamma function

THINGS TO TRY:


- gamma function
- gamma function of 2
- gamma function of (1/2)

Interactive knowledge apps from
Wolfram Demonstrations Project

Graphs of the Beta Function
Daniel de Souza Carvalho

Continuous Binomial Function in 3D
Larry Keasler

Fractional Derivative of a Power
Jorge Gamaliel Frade Chávez

COMPUTE &
VISUALIZE
JUST ABOUT ANYTHING

Try Mathematica
for Free
Desktop + Online

a slightly unfortunate notation due to Legendre which is now universally used instead of Gauss's simpler $\Pi(n) = n!$ (Gauss 1812; Edwards 2001, p. 8).

It is analytic everywhere except at $z = 0, -1, -2, \dots$, and the residue at $z = -k$ is

$$\text{Res } \Gamma(z) = \frac{(-1)^k}{k!}.$$
(2)

There are no points z at which $\Gamma(z) = 0$.

The gamma function is implemented in the Wolfram Language as `Gamma[z]`.

There are a number of notational conventions in common use for indication of a power of a gamma functions. While authors such as Watson (1939) use $\Gamma^x(z)$ (i.e., using a trigonometric function-like convention), it is also common to write $[\Gamma(z)]^x$.

The gamma function can be defined as a definite integral for $R(z) > 0$ (Euler's integral form)


$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt$$
(3)

$$= 2 \int_0^\infty e^{-t^2} t^{z-1} dt.$$
(4)

or

$$\Gamma(z) = \int_0^\infty \left[\ln\left(\frac{1}{t}\right) \right]^{z-1} dt.$$
(5)

The complete gamma function $\Gamma(x)$ can be generalized to the upper incomplete gamma function $\Gamma(a, x)$ and lower incomplete gamma function $\gamma(a, x)$.



Plots of the real and imaginary parts of $\Gamma(z)$ in the complex plane are illustrated above.

Integrating equation (3) by parts for a real argument, it can be seen that

Algebra

Applied Mathematics

Calculus and Analysis

Discrete Mathematics

Foundations of Mathematics

Geometry

History and Terminology

Number Theory

Probability and Statistics

Recreational Mathematics

Topology

Alphabetical Index

Interactive Entries

Random Entry

New in MathWorld

MathWorld Classroom

About MathWorld

Contribute to MathWorld

Send a Message to the Team

MathWorld Book

Wolfram Web Resources

13,612 entries

Last updated: Mon Feb 27 2017

Created, developed, and
nurtured by Eric Weisstein
at Wolfram Research

Calculus and Analysis > Special Functions > Gamma Functions >
Calculus and Analysis > Series > Series Expansions >

Stirling's formula for n!

Stirling's Approximation

DO DOWNLOAD
WolframAlpha

CONTRIBUTE
TO THIS PAGE

Stirling's approximation gives an approximate value for the factorial function $n!$ or the gamma function $\Gamma(n)$ for $n \gg 1$. The approximation can most simply be derived for n an integer by approximating the sum over the terms of the factorial with an integral, so that

$$\ln n! = \ln 1 + \ln 2 + \dots + \ln n \quad (1)$$

$$= \sum_{k=1}^n \ln k \quad (2)$$

$$\approx \int_1^n \ln x \, dx \quad (3)$$

$$= [x \ln x - x]_1^n \quad (4)$$

$$= n \ln n - n + 1 \quad (5)$$

$$\approx n \ln n - n. \quad (6)$$

The equation can also be derived using the integral definition of the factorial,

$$n! = \int_0^\infty e^{-x} x^n \, dx. \quad (7)$$

Note that the derivative of the logarithm of the integrand can be written

$$\frac{d}{dx} \ln(e^{-x} x^n) = \frac{d}{dx} (n \ln x - x) = \frac{n}{x} - 1. \quad (8)$$

The integrand is sharply peaked with the contribution important only near $x = n$. Therefore, let $x \equiv n + \xi$ where $\xi \ll n$, and write

$$\ln(x^n e^{-x}) = n \ln x - x \quad (9)$$

$$= n \ln(n + \xi) - (n + \xi). \quad (10)$$

Now,

$$\ln(n + \xi) = \ln\left[n\left(1 + \frac{\xi}{n}\right)\right] \quad (11)$$

$$= \ln n + \ln\left(1 + \frac{\xi}{n}\right) \quad (12)$$

$$= \ln n + \frac{\xi}{n} - \frac{1}{2} \frac{\xi^2}{n^2} + \dots \quad (13)$$

so

$$\ln(x^n e^{-x}) = n \ln(n + \xi) - (n + \xi) \quad (14)$$

$$= n \ln n + \xi - \frac{1}{2} \frac{\xi^2}{n} - n - \xi + \dots \quad (15)$$

$$= n \ln n - n - \frac{\xi^2}{2n} + \dots \quad (16)$$

Taking the exponential of each side then gives

$$x^n e^{-x} \approx e^{\ln n} e^{-n} e^{-\xi^2/2n} \quad (17)$$

$$= n^n e^{-n} e^{-\xi^2/2n}. \quad (18)$$

Plugging into the integral expression for $n!$ then gives

$$n! \approx \int_{-n}^n n^n e^{-n} e^{-\xi^2/2n} d\xi \quad (19)$$

$$\approx n^n e^{-n} \int_{-\infty}^{\infty} e^{-\xi^2/2n} d\xi. \quad (20)$$

Evaluating the integral gives

$$n! \approx n^n e^{-n} \sqrt{2\pi n} \quad (21)$$

$$= \sqrt{2\pi} n^{n+1/2} e^{-n}. \quad (22)$$

(Wells 1986, p. 45). Taking the logarithm of both sides then gives

$$\ln n! \approx n \ln n - n + \frac{1}{2} \ln(2\pi n) \quad (23)$$

$$= \left(n + \frac{1}{2}\right) \ln n - n + \frac{1}{2} \ln(2\pi). \quad (24)$$

This is Stirling's series with only the first term retained and, for large n , it reduces to Stirling's approximation

$$\ln n! \approx n \ln n - n. \quad (25)$$


Taking successive terms of $\lfloor n^k/n! \rfloor$, where $\lfloor x \rfloor$ is the floor function, gives the sequence 1, 2, 4, 10, 26, 64, 163, 416, 1067, 2755, ... (OEIS A005775).

Stirling's approximation can be extended to the double inequality

$$\sqrt{2\pi} n^{n+1/2} e^{-n+1/(12(n+1))} < n! < \sqrt{2\pi} n^{n+1/2} e^{-n-1/(12n)}. \quad (26)$$

(Robbins 1955, Feller 1968).

Gosper has noted that a better approximation to $n!$ (i.e., one which approximates the terms in Stirling's series instead of truncating them) is given by



P1.4

$$\int \frac{1}{1+\sin^2 x} dx = \int \frac{y = \tan x}{\sin^2 x = \frac{\tan^2 x}{\sin^2 x}} \frac{\sqrt{1+\frac{\pi}{2}+k^2} dy}{\sin^2 x + \cos^2 x} \quad \left. \begin{array}{l} dy = \frac{dx}{\cos^2 x} \\ \frac{dy}{1+y^2} = dx \end{array} \right\} (P+2)$$

cada $\sin x$ a $\tan x$

$$= 4x \int \frac{1}{1+\sin^2 x} \frac{1}{\frac{1}{\frac{1}{2}}} dx$$

$$\sqrt{\frac{\pi}{2}} = \frac{2\sqrt{2}\pi}{4}$$


$$= \int \frac{1}{1+\frac{z^2}{1+y^2}} \frac{dy}{1+y^2}$$

$$\cos^2 x = \frac{\cos^2 x}{\cos^2 x + \sin^2 x}$$

$$= \frac{1}{1+\tan^2 x}$$

$$= \frac{1}{1+y^2}$$

$$\frac{1}{1+\sin^2 x}$$



$$= \int \frac{1}{1+z^2} \frac{dz}{1+y^2} = \int \frac{1}{1+z^2} dz$$

$$\left. \begin{array}{l} \bar{z}^2 y = z \\ \bar{z}^2 dy = dz \end{array} \right\} = \frac{1}{\bar{z}^2} \int \frac{1}{1+z^2} dz = \frac{1}{\bar{z}^2} \arctan z$$

$$= \frac{1}{\bar{z}^2} \arctan(\bar{z}^2 y) = \frac{1}{\bar{z}^2} \arctan(\bar{z}^2 \tan x) = F(x)$$


• Frage: $\int_{\alpha}^{\pi} \frac{1}{1+\sin^2 x} dx = F(\pi) - F(\alpha) = 0?$

Newton-Leibniz


f(x) stetig > 0
auf $(\alpha, \bar{\alpha})$

$$f(x) = \frac{1}{1+\sin^2 x}$$

$$F(x) = \frac{1}{\bar{z}^2} \arctan(\bar{z}^2 \tan x) \text{ auf } \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$$



$$\lim_{x \rightarrow \frac{\pi}{2}^-} F(x) = \lim_{x \rightarrow \frac{\pi}{2}^+} \frac{1}{\bar{z}^2} \arctan(\bar{z}^2 \tan x) = \pm \frac{\pi}{\bar{z}^2}$$



$$F(x) \text{ bei } x \in \{\alpha, \bar{\alpha}\}$$

$$\tilde{F}(x) \text{ bei } x \in \{\alpha, \bar{\alpha}\} \quad \left\{ \begin{array}{l} \frac{\pi}{\bar{z}^2} \quad x = \frac{\pi}{2} \\ F(x) + \frac{\pi}{\bar{z}^2} \quad x \in (\frac{\pi}{2}, \bar{\alpha}) \end{array} \right.$$

$\int \frac{1}{\sin^4 x + \cos^4 x} dx = \frac{\pi}{2}$ P1-3
 $= \frac{1}{2} \cdot \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}} \frac{1}{\sin^4 x + \cos^4 x} dx$
 $I_4 = \int_{-\infty}^{\infty} \frac{1}{(1+y^2)^2 + (1-y^2)^2} dy = \int_{-\infty}^{\infty} \frac{1}{4y^2} dy$
 $= \int_{-\infty}^{\infty} \frac{1+iy^2}{1+iy^2} dy$
 $\Rightarrow z = \pm \frac{\pi}{2} (\pm i \pm 1)$
 $\Rightarrow (z - \frac{\pi}{2}(1+i))(z - \frac{\pi}{2}(1-i))(z + \frac{\pi}{2}(1+i))(z + \frac{\pi}{2}(1-i))$
 $= (z^2 - \frac{\pi^2}{2}z + 1)(z^2 + \frac{\pi^2}{2}z + 1)$
 $= \frac{1}{2} \left(\int_{-\infty}^{\infty} \frac{1}{z^2 + \frac{\pi^2}{2}z + 1} + \int_{-\infty}^{\infty} \frac{1}{z^2 - \frac{\pi^2}{2}z + 1} \right) dz$
 $= \frac{i}{2} \left(\int_{-\infty}^{\infty} \frac{dz}{(z + \frac{\pi}{2})^2 + \frac{1}{4}} + \frac{i}{2} \left(\int_{-\infty}^{\infty} \frac{dz}{(\frac{z}{2} - \frac{\pi}{2})^2 + \frac{1}{4}} \right) \right)$
 $= \frac{i}{2} \left(\int_{-\infty}^{\infty} \frac{dz}{(\frac{z}{2} + \frac{\pi}{2})^2 + 1} + \frac{i}{2} \left(\int_{-\infty}^{\infty} \frac{dz}{(\frac{z}{2} - \frac{\pi}{2})^2 + 1} \right) \right)$
 $= \frac{1}{2} \left(\int_{-\infty}^{\infty} \frac{dz}{z^2 + 1} + \int_{-\infty}^{\infty} \frac{dz}{z^2 + 1} \right)$
 $= \frac{1}{2} [\arctan(z) + \arctan(-z)]$
 $= \frac{1}{2} [\arctan(\frac{z}{2} + \frac{\pi}{2}) + \arctan(\frac{z}{2} - \frac{\pi}{2})]$
 $I_4 = \int_{-\infty}^{\infty} \frac{1}{\sin^4 x + \cos^4 x} dx = \dots \frac{\pi}{2}$
 $= \frac{1}{2} \left[\frac{\pi}{2} - \arctan(1) \right] = \frac{\pi}{2} + \frac{\pi}{2} - \arctan(-1)$

$$\text{P1.6} \quad \int_{\frac{\pi}{2}}^{\infty} \frac{1}{x^2} dx = -\frac{1}{x} \Big|_{\frac{\pi}{2}}^{\infty} = 0 - \left(-\frac{1}{\frac{\pi}{2}}\right) = \frac{2}{\pi}$$

$$\text{P1.7} \quad \int_0^{\infty} e^{3x} dx = \left[-\frac{e^{-3x}}{3} \right]_0^{\infty} = -0 + \frac{1}{3} = \frac{1}{3}$$

$$\text{P1.8} \quad \int x^2 \ln x dx = \begin{cases} \text{mehr } n=2 \\ n=x \quad n=\frac{1}{2} \end{cases} = \left[\frac{1}{2} x^2 \ln x \right]_1^1 - \frac{1}{2} \int x^2 dx$$

$$= \left[-\frac{x^3}{9} \right]_1^1 = -\frac{1}{9}$$

\$e^{ix} = \cos bx + i \sin bx\$

$$\text{P1.9} \quad \int e^{ax} \cos(bx) dx = \operatorname{Re} \int e^{(a+ib)x} dx$$

$$= \operatorname{Re} \left(\frac{e^{(a+ib)x}}{-a+ib} \right) = \operatorname{Re} \left(e^{-ax} \frac{e^{ibx}}{-a+ib} \right)$$

$$= \operatorname{Re} \left(e^{-ax} \frac{\cos bx + i \sin bx}{-a+ib} \right) = \operatorname{Re} \left(e^{-ax} \frac{\cos bx + i \sin bx ((a-i)b)}{a^2 + b^2} \right)$$

$$= \frac{e^{-ax}}{a^2 + b^2} \left(-a \cos bx + b \sin bx \right) \Big|_0^{\infty} = \frac{0 - a}{a^2 + b^2} \quad (+?)$$

$$\text{P1.10} \quad \int_{\frac{\pi}{2}}^{\infty} \tan x dx$$

$$\int \tan x dx = \int \frac{\sin x}{\cos x} dx \left| \begin{array}{l} t = \cos x \\ dt = -\sin x dx \end{array} \right. = - \int \frac{dt}{t}$$

$$= -\ln|t| = -\ln(\cos x)$$

$$\rightarrow \approx -\ln(\cos x)^{\frac{1}{2}}, \quad \text{= unlogar.}$$

Fr. 11

$$\int_0^{\pi} \ln(1+2x\cos x + a^2) dx$$

(Fr. 5)

$$z = \pi - x$$

$$dz = -dx$$

$$x=0 \rightarrow z=\pi$$

$$x=\pi \rightarrow z=0$$

$$\begin{aligned} & \left. \int_0^{\pi} \ln(1+2x\cos x + a^2) dx \right. \\ & \quad - \int_{\pi}^0 \ln(1+2z\cos z + a^2) dz \\ & = \int_0^{\pi} \ln(1+2z\cos z + a^2) dz \end{aligned}$$

$$\cos(\pi-z) =$$

$$-\cos z$$

$$I(a) = I(-a)$$

$$I(a) + I(-a) = \int_0^{\pi} (\ln(1+2x\cos x + a^2) + \ln(1-2x\cos x + a^2)) dx$$

$$= \int_0^{\pi} \ln((1+2a\cos x + a^2)(1-2a\cos x + a^2)) dx$$

$$= \int_0^{\pi} \ln((1+a^2)^2 - (2a\cos x)^2) dx$$

$$\cos^2(2x) = \cos^2 x - \sin^2 x \\ = 2\cos^2 x - 1$$

$$= \int_0^{\pi} \ln(1+2a^2 + a^4 - 2a^2(1+\cos 2x)) dx$$

$$= \int_0^{\pi} \ln(1-2a^2 \cos 2x + a^4) dx$$

$$= \left\{ \begin{array}{l} y = 2x \\ dy = 2dx \\ y = 0, \dots, 2\pi \end{array} \right. = \frac{1}{2} \int_0^{2\pi} \ln(1-2a^2 \cos y + a^4) dy$$

$$= \frac{1}{2} \int_0^{\pi} \ln(1-2a^2 \cos y + a^4) dy + \frac{1}{2} \int_{\pi}^{2\pi} \ln(1-2a^2 \cos y + a^4) dy$$

$$= \frac{1}{2} \int_0^{\pi} \ln(1-2a^2 \cos y + a^4) dy + \left\{ \begin{array}{l} x = 2\pi - y \\ dx = -dy \\ x = \pi, \dots, 0 \end{array} \right\} - \int_{\pi}^0 \ln(1-2a^2 \cos x + a^4) dx$$

$$= \frac{1}{2} I(a^2) + \frac{1}{2} I(a^2) = I(a^2)$$

$$2I(a) = I(a^2)$$

$$a=0 \quad \cancel{2I(0)} \quad 2I(0) = I(0) \Rightarrow I(0)=0$$

$$a=1 \quad 2I(1) = I(1) \Rightarrow I(1)=0$$

For $|a| < 1$

$$I(a) = \frac{1}{2} I(a^2) = \frac{1}{2} \cdot \frac{1}{2} I(a^4) = \dots = \frac{1}{2^n} I(a^{2n})$$


$$\text{as } n \rightarrow \infty \quad I(a) = \lim_{n \rightarrow \infty} \frac{1}{2^n} I(a^{2n}) = 0$$

For $a > 1$

$$\begin{aligned} I(a) &= \int_0^\pi \ln \left\{ a^2 \left(\frac{1}{a^2} + \frac{2}{a} \cos x + 1 \right) \right\} dx \\ &= \underbrace{\int_0^\pi \ln a^2 dx}_{= 2 \ln a} + \underbrace{\int_0^\pi \ln \left(1 + \frac{2}{a} \cos x + \frac{1}{a^2} \right) dx}_{I(\gamma_a) = 0} \\ &= 2 \ln a \end{aligned}$$

$$\Rightarrow I(a) = \begin{cases} 0 & \text{if } |a| \leq 1 \\ 2 \ln a & \text{if } |a| > 1 \end{cases}$$

Fazehlini mukave



$$\oint \vec{E} \cdot d\vec{s} = 2\pi r E(r) \hat{x} = \frac{dN}{dx} = \frac{dk}{dx}$$

$$\vec{E} = \frac{x}{2\pi r c} \hat{e}_x$$

$$\vec{E} = -\nabla V$$

$$V = -\frac{x}{2\pi r c} \ln r$$

Line
Dose

$$\text{Line} \rightarrow \text{cylinder} \rightarrow r \rightarrow \ln(1 - 2r \cos \theta + r^2) \rightarrow \frac{1}{2} \ln(1 - 2r \cos \theta + r^2)$$

Riemann sum

P1-7

$$\int \ln(1 - 2a \cos x + a^2) dx$$

$$(a-z)(a-z^*)$$

$$= (a-(b+ic))(a-(b-ic))$$

$$= a^2 - a(b+ic + b-ic) + 2\ln(b+ic)$$

$$= \frac{\pi}{m} \left\{ \ln \left(1 - 2a \cos \frac{\pi}{m} + a^2 \right) + \ln \left(1 - 2a \cos \frac{2\pi}{m} + a^2 \right) + \dots + \ln \left(1 - 2a \cos \frac{(m-1)\pi}{m} + a^2 \right) \right\}$$

$$= \frac{\pi}{m} \ln \prod_{k=1}^m \left(1 - 2a \cos \left(\frac{k\pi}{m} \right) + a^2 \right)$$

$$z = b+ic$$

$$(a-z)(a-z^*) = a^2 - a(z+z^*) + zz^*$$

$$= a^2 + a(b+ic+b-ic) + (b+ic)(b-ic)$$

$$= a^2 - 2ab + b^2 + c^2$$

$$\begin{matrix} \uparrow & \uparrow & \uparrow \\ \cos(k) & \cos^2 + \sin^2 & z = \cos + i \sin \end{matrix}$$

$$= \frac{\pi}{m} \ln \prod_{k=1}^m \left[a - \left(\cos \frac{k\pi}{m} + i \sin \frac{k\pi}{m} \right) \right] \left[a - \left(\cos \left(\frac{k\pi}{m} \right) + i \sin \left(\frac{k\pi}{m} \right) \right) \right] (a \neq 1)$$

~~a different~~

$$(a+1)(a-1)$$

$$a^{2m} - 1$$

$$a^{2m} = 1$$

$$= \frac{\pi}{m} \ln \frac{1-a^{2m}}{1-a^2} = \frac{\pi}{m} \ln \frac{1-(a^2)^m}{1-a^2}$$

$$\text{if } |a| < 1 \quad \lim_{m \rightarrow \infty} \frac{\pi}{m} \ln \ln \frac{1-a^{2m}}{1-a^2} = 0$$

$$\text{if } |a| > 1 \quad \lim_{m \rightarrow \infty} \frac{\pi}{m} \ln \frac{1-a^{2m}}{1-a^2} = \lim_{m \rightarrow \infty} \frac{\pi}{m} \ln \frac{a^{2m}-1}{a^2-1}$$

$$= \lim \left\{ \frac{\pi}{m} \ln (a^{2m}-1) - \ln (a^2-1) \right\}$$

$$= \lim_{m \rightarrow \infty} \left(\frac{\pi}{m} 2ma - \underbrace{\frac{\pi}{m} \ln(a^2-1)}_{\ln(a^2-1)} \right)$$

$$= 2\pi \ln a$$

P 1.12

$$\int_a^{\infty} x^p dx \quad p \in \mathbb{R}$$

Plaus

$$p \neq -1 \quad \int x^p = \frac{x^{p+1}}{p+1}$$

$$p = -1 \quad (\star) \frac{1}{x} = \ln x$$

$$\int_a^{\infty} x^p = \lim_{x \rightarrow \infty} \frac{x^{p+1}}{p+1} - \lim_{x \rightarrow a} \frac{x^{p+1}}{p+1}$$

$$\int_a^{\infty} \frac{1}{x} = \lim_{x \rightarrow \infty} \ln x - (\lim_{x \rightarrow a} \ln x + \infty - \infty)$$

$$\rightarrow p > -1 : \lim_{x \rightarrow \infty} \frac{x^p}{x} - \lim_{x \rightarrow a^+} \frac{x^p}{x} = \infty - 0$$

$$p < -1 : -\lim_{x \rightarrow \infty} \frac{1}{p x^p} + \lim_{x \rightarrow a^+} \frac{1}{p x^p} = -\infty + \infty$$

Integral nicht reellwertig.

P 1.13 $\int_a^{\infty} x^p dx \quad p \in \mathbb{R}$: $p < -1 \quad \int_a^{\infty} x^p dx = -\frac{1}{p+1} \checkmark$
 Laut Kopáček ist dies falsch.

P 1.14 $\int_a^{\infty} x^p dx \quad p \in \mathbb{R}$: $p > -1 \quad \int_a^{\infty} x^p dx = \frac{1}{p+1} \checkmark$

P 1.15 $\int_a^{\infty} \frac{x^{3/2}}{1+x^2} dx = \int_a^{\infty} \frac{x^{3/2}}{1+x^2} dx + \int_a^{\infty} \frac{x^{3/2}}{1+x^2} dx$

Sei $Kopáček$, p. 155 wichen

$$x \rightarrow 0 : \frac{x^{3/2}}{1+x^2} \sim x^{3/2} \quad \text{dann}$$

$$x \rightarrow \infty : \frac{x^{3/2}}{1+x^2} \sim \frac{1}{x^{1/2}} \quad \lim_{x \rightarrow \infty} \frac{x^{3/2}}{1+x^2} \sim \frac{x^{3/2}}{x^{3/2}} = \lim_{x \rightarrow \infty} \frac{1}{1+\frac{1}{x^2}} = 1$$

für spezielle reellwertige Funktionen auf (a, ∞)

$\lim_{x \rightarrow a^+} \frac{f}{g} = \text{feste reellwertige} \rightarrow \text{da lösbar, also da reellwertig}$

$$\int_a^{\infty} f(x) dx \quad \int_a^{\infty} g(x) dx$$

$$\rightarrow x^p \text{ für } p = -\frac{1}{2} \rightarrow \text{unendlich}$$

Dafür gilt $p < -1$

$$\bullet \int_0^{\infty} \frac{x^{3/2}}{1+x^2} dx \leq \int_0^{\infty} \frac{x^{3/2}}{x^2} dx = \left[x^{-1/2} \right]_0^{\infty} = 2$$

(77-9)

$$\bullet \int_0^{\infty} \frac{x^{3/2}}{1+x^2} \geq \int_0^{\infty} \frac{x^{3/2}}{2x^2} dx = 2 \left[x^{-1/2} \right]_0^{\infty} = \left[4x^{-1/2} \right]_0^{\infty} = \infty$$

Nehmung (noch zu klären)

P1.16 $\int_0^{\infty} \frac{1}{\sqrt{t-x^2}} dx$

$f(x) > 0 \text{ für } (0, t)$

$$\stackrel{x \rightarrow 0+}{\sim} \frac{1}{\sqrt{t-x^2}} \sim x^{-1/2} \quad p > -1 \quad \text{OK?}$$

$$\stackrel{x \rightarrow t-}{\sim} \frac{1}{\sqrt{t-x^2}} \sim \frac{1}{\sqrt{(t+x)(t-x)}} \sim \frac{1}{\sqrt{t-x}}$$

$$\int_0^{\infty} \frac{dx}{\sqrt{t-x^2}} \leq \int_0^{\infty} \frac{1}{\sqrt{t}} = \frac{1}{\sqrt{t}} \left[x \right]_0^{\infty}$$

$$\int_{\pi/2}^{\pi/2} \frac{dx}{\sqrt{t-x^2}} \leq \sqrt{t} \int_{\pi/2}^{\pi/2} \frac{dx}{\sqrt{t-x^2}} = \sqrt{t} \left[\arcsin x \right]_{\pi/2}^{\pi/2} = \sqrt{t} \left(\frac{\pi}{2} - \frac{\pi}{2} \right) = 0$$

P1.17 $\int_{-\infty}^{\infty} \frac{1}{e^{xt}} dx$ $\begin{cases} x=t \\ dx=e^t dt \end{cases} \quad \begin{cases} \int_{-\infty}^{\infty} \frac{e^t}{t} dt \\ = \int_{-\infty}^{\infty} \frac{e^t}{t} dt \end{cases}$

$$\int_{-\infty}^{\infty} \frac{e^t}{t} dt \geq \int_{-\infty}^{\infty} \frac{1}{t} dt = \lim_{a \rightarrow \infty} \left[\ln|t| \right]_{-a}^a = \lim_{a \rightarrow \infty} (\ln a - (-a)) = \infty$$

$$\int_{-\infty}^{\infty} \frac{e^t}{t} dt = \int_{-\infty}^{\infty} \frac{e^t}{t} dt + \int_{-\infty}^{\infty} \frac{e^t}{t} dt \leq \int_{-\infty}^{\infty} \frac{e^t}{t} dt + \int_{-\infty}^{\infty} \frac{e^t}{t} dt = 2 \int_{-\infty}^{\infty} \frac{e^t}{t} dt = -\infty$$

\rightarrow nicht konvergiert X

P1.18

$$\int_0^{\pi/2} \frac{\ln \sin x}{x^p} dx \quad p \in \mathbb{R}$$

$p=0$ ✓
 $p=1$ ✗
 $0 < p < 1$ ✓

$$\begin{aligned} \lim_{x \rightarrow 0^+} \frac{\ln \sin x}{x^p} &= \begin{cases} -\infty & p > 0 \\ \text{D.L.H.} & p = 0 \\ \lim_{x \rightarrow 0^+} \frac{\cos x}{p x^{p-1}} & 0 < p < 1 \end{cases} \sim \frac{1}{p} \lim_{x \rightarrow 0^+} \frac{\cos x}{\sin x} \frac{x^{1-p}}{x^p} \\ &= \frac{1}{p} \lim_{x \rightarrow 0^+} \left(\frac{\cos x}{\sin x} \right) \frac{x^{-p}}{1} = 0 \quad \text{pro } p < 0 \end{aligned}$$

$p \geq 0$

$$\begin{aligned} \int_0^{\pi/2} \frac{\ln \sin x}{x^p} dx &\leq \int_0^{\pi/2} \frac{\ln x}{x^p} dx \quad \left\{ \begin{array}{l} u = \ln x \quad u' = \frac{1}{x} \\ v = x^{-p} \quad v' = \frac{x^{1-p}}{1-p} \end{array} \right\} \\ &= \left[\ln x \frac{x^{1-p}}{1-p} \right]_0^{\pi/2} - \int_0^{\pi/2} \frac{x^{-p}}{1-p} dx \\ &= \left[\ln x \frac{x^{1-p}}{1-p} - \frac{x^{1-p}}{(1-p)^2} \right]_0^{\pi/2} = \frac{1}{(1-p)^2} \left[x^{1-p} \left((1-p) \ln x - 1 \right) \right]_0^{\pi/2} \end{aligned}$$

* $p > 1 \Rightarrow -\infty$

* $p = 1 \Rightarrow$

P1.15

$$\int_0^{\infty} \frac{\arctan x}{x^{3/2}} dx$$

$$\frac{\arctan x}{x^{3/2}} > 0 \text{ na } (0, \infty)$$

$$\int_0^1 \frac{\arctan x}{x^{3/2}} dx \leq \underbrace{\int_0^1 \frac{x}{x^{3/2}} dx}_{\arctan x \leq x} = \int_0^1 \frac{dx}{\sqrt{x}} = 2 [\sqrt{x}]_0^1 = 2$$

$$\int_1^{\infty} \frac{\arctan x}{x^{3/2}} dx \leq \frac{\pi}{2} \int_1^{\infty} \frac{1}{x^{3/2}} dx = \frac{\pi}{2} \left[x^{-1/2} \right]_1^{\infty} = \frac{\pi}{2} \left(\frac{1}{-\frac{1}{2}} \right)_1^{\infty}$$

$$= \pi \left[\frac{1}{\sqrt{x}} \right]_{\infty}^1 = \pi$$

(R)

Věta (7.6.5) Srovnání vlnic pro brouzgou? Nejdovorejší funkce

Nedle $-\infty < a < b \leq +\infty$

• Testovací $f \in C([a,b])$, $g \geq 0$ neobsahující nula na (a,b) a $f(x) = O(g(x))$ pro $x \rightarrow b_-$ \Rightarrow pak i f je neobsahující nula na (a,b)

• Testovací $f,g \in C([a,b])$ nezáporné, $f(x) = O(g(x)) \Leftrightarrow g(x) = O(f(x))$ pro $x \rightarrow b_-$ \Rightarrow pak f neutr. integr. na (a,b) $\Leftrightarrow g$ neutr. integr. na (a,b)

• Speciálně pokud $f,g,h \in C([a,b])$ nezáporné, $f(x) = O(g(x)) \circ g(x) = O(h(x))$ pro $x \rightarrow b_-$, pak fh je neobsahující nula na (a,b) $\Leftrightarrow gh$ neutr. integr. na (a,b)

Příklad (N) $\int_0^1 \frac{e^x - 1}{x \sin x} dx$ $g = \frac{1}{x}$ $\lim_{x \rightarrow 0^+} \frac{f(x)}{g(x)} = \lim_{x \rightarrow 0^+} \frac{e^x - 1}{\sin x} = 1$ (N) $\int_0^1 \frac{1}{x} dx = [2\ln x]_0^1 = 2 \in \mathbb{R}$

Příklad (N) $\int_0^1 x \ln x \sin \frac{1}{x} dx$ $g = |\ln x| = -\ln x$ $|f| \leq g$ na $(0,1)$
 $(N) \int_0^1 -\ln x dx = -[x \ln x - x]_0^1 = 1 \Rightarrow \int_0^1 x \ln x \sin \frac{1}{x} dx$ existuje

Příklad (N) $\int_0^{+\infty} \arctan x dx$ $g(x) = 1$... neutr. integr., zároveň $\lim_{x \rightarrow +\infty} \frac{f(x)}{g(x)} = \frac{\pi}{2} \Rightarrow$
 \Rightarrow existuje

Věta (7.6.11) Abel-Dirichletův kritérium

Nedle $F, f, g \in C([a,b])$, $F' = f$ na (a,b) a g monotonní na (a,b)

• (Dirichlet) Je-li F mezená na (a,b) a $\lim_{x \rightarrow b_-} g(x) = 0 \Rightarrow fg$ neutr. integr. na (a,b)

• (Abel) Je-li f neutr. integr. na (a,b) a g mezená na (a,b) $\Rightarrow fg$ neutr. integr. na (a,b)

Příklad: $\int_a^\infty \frac{\sin x}{x} dx$ $f = \sin x$... $F = -\cos x$ mezená
 $g = \frac{1}{x}$... monotonní, $\lim_{x \rightarrow +\infty} \frac{1}{x} = 0$ } $\Rightarrow \int_a^\infty \frac{\sin x}{x} dx$ existuje

$$\int_a^\infty \frac{\sin x}{x} dx = \int_a^\infty \left(\frac{1}{2x} - \frac{\cos(2x)}{2x} \right) dx$$

↗ kde Dirichlet

$$f = \sin x \dots F \text{ mezená} \left(= \frac{x}{2} - \frac{1}{2} \sin(2x) \right)$$